



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

or, $\frac{h(\sin C \sin E - \cos C \sin E)}{y(\sin A \cos E + \cos A \sin E)} = \frac{\sin C}{\sin A}$; or $h(1 - \cot C \tan E) = y(1 + \cot A \tan E)$, or

$$h - y = (h \cot C + y \cot A) \tan E; \text{ but } \tan E = (h - y)/x,$$


$\therefore x = y \cot A + h \cot C$, or $y = x \tan A - h \cot C \tan A$, as the locus of C .

For $C = A$, we have $y = x \tan A - h$.

Make $OF = OA$, draw FH so as to make $\angle HGN = \angle A$, draw AG . $\angle HGN = \angle OGF = \angle AGO = \angle A$.

Since $\angle BAC = \angle BGC$, $ABCG$ is concyclic.

$\therefore \angle ACB = \angle AGB = \angle A$, $\therefore \angle C = \angle A$, which unifies geometrically for the case in which the angles at A and C remain equal.



92. Proposed by JOSIAH H. DRUMMOND, LL. D., Counselor at Law, Portland, Me.

Let $ABCD$ be a quadrilateral inscribed in a circle. Draw the diagonals AC and BD . Show that $AB \cdot BC : DC \cdot AD = BD : AC$. [From a note in *Young's Geometry*, edition of 1830.]

Solution by the PROPOSER, and J. SCHEFFER, Hagerstown, Md.

Let F be the intersection of the diagonals.

Then $AB : BF :: CD : CF$,

or $AB : CD :: BF : CF$,

and $BC : AD :: CF : DF$.

Hence $AB \cdot BC : AD \cdot CD :: BF : DF$,

(I) and $AB \cdot BC + AD \cdot CD : AD \cdot CD :: BF + DF (= BD) : DF$.

In like manner it is shown

(II) that $AB \cdot AD + BC \cdot CD : BC \cdot DC :: AC : CF$.

But $AD : DF :: BC : CF$,

or $AD \cdot CD : DF :: BC \cdot CD : CF$.

Combining these with (I) and (II), we have

$$AB \cdot BC + AD \cdot CD : AB \cdot AD + BC \cdot CD :: BD : AC. \quad \text{Q. E. D.}$$

Also solved by B. F. SINE, CHAS. C. CROSS, WALTER H. DRANE, and G. B. M. ZERR.

93. Proposed by G. B. M. ZERR, A. M., Ph. D., President and Professor of Mathematics, The Russell College, Lebanon, Va.

While surveying in a level field I notice a mountain behind a hill. Wish-
ing to know the height of each I take the angles of elevation of the tops of both
and find them to $\beta = 45^\circ$, $\delta = 40^\circ$. I then measure a straight line $a = 400$ feet,
and find the angles of elevation of the tops to be $\gamma = 42^\circ$, $\mu = 38^\circ$. After measur-
ing $b = 300$ feet more in the same straight line I find the elevations to be $\lambda = 40^\circ$,
 $\nu = 36^\circ$. Find the height of each.

Solution by the PROPOSER.

Let $AB = 400$ feet $= a$, $BC = 300$ feet $= b$, $OP = x$, $QR = y$.

$\angle OAP=45^\circ=\beta$, $\angle OAQ=40^\circ=\delta$, $\angle OBP=42^\circ=\gamma$, $\angle OBQ=38^\circ=\mu$,
 $\angle OCP=40^\circ=\lambda$, $\angle OCQ=36^\circ=\nu$.

$OA=m$, $OB=n$, $OC=p$, $\angle OCA=b$.

$\therefore m=y \cot \delta = x \cot \beta$, $n=y \cot \mu = x \cot \gamma$.


$p=y \cot \nu = x \cot \lambda$.

Also from triangles OCA and OCB ,

$$m^2 = p^2 + (a+b)^2 - 2p(a+b) \cos \theta \dots\dots\dots (1).$$

$$n^2 = p^2 + b^2 - 2pb \cos \theta \dots\dots\dots (2).$$

The values of m , n , p in (1) and (2) and eliminating $\cos \theta$, we get



$$x = \sqrt{\frac{ab(a+b)}{b \cot^2 \beta + a \cot^2 \lambda - (a+b) \cot^2 \gamma}}$$

$$y = \sqrt{\frac{ab(a+b)}{b \cot^2 \delta + a \cot^2 \nu - (a+b) \cot^2 \mu}}$$

Substituting we get $y=1505.183$ feet, $x=4232.505$ feet.

Also solved by *ALOIS F. KAVORIK*, and *CHAS. C. CROSS*.

CALCULUS.

71. Proposed by *J. C. CORBIN*, Pine Bluff, Ark.

Form the differential equation of the third order, of which

$y=c_1 e^{2x}+c_2 e^{-3x}+c_3 e^x$ is the complete primitive.

III. Solution by *C. HORNING*, A. M., Professor of Mathematics, Heidelberg University, Tiffin, O.; *COOPER D. SCHMITT*, A. M., Professor of Mathematics, University of Tennessee, Knoxville, Tenn.; *C. W. M. BLACK*, A. M., Professor of Mathematics, Wesleyan Academy, Wilbraham, Mass.; *G. B. M. ZERR*, A. M., Ph. D., The Russell College, Lebanon, Va.; and the PROPOSER.

$$y=c_1 e^{2x}+c_2 e^{-3x}+c_3 e^x \dots\dots (1), \quad dy/dx=2c_1 e^{2x}-3c_2 e^{-3x}+c_3 e^x \dots\dots (2);$$

$$d^2 y/dx^2=4c_1 e^{2x}+9c_2 e^{-3x}+c_3 e^x \dots\dots (3). \quad d^3 y/dx^3=8c_1 e^{2x}-27c_2 e^{-3x}+c_3 e^x \dots\dots (4).$$

$$(2)-(1) \text{ gives } dy/dx-y=c_1 e^{2x}-4c_2 e^{-3x} \dots\dots\dots (5),$$

$$(4)-(1) \text{ gives } d^3 y/dx^3-y=7c_1 e^{2x}-28c_2 e^{-3x} \dots\dots\dots (6),$$

$$(6)-7(5) \text{ gives } d^3 y/dx^3-7(dy/dx)+6y=0.$$

IV. Solution by *J. SCHEFFER*, A. M., Hagerstown, Md.

The equation reduces itself to what is the equation the three roots of which are 1, 2, -3 . This equation is $z^2-7z+6=0$; consequently the complete primitive is $d^3 y/dx^3-7(dy/dx)+6y=0$.

72. Proposed by *G. B. M. ZERR*, A. M., Ph. D., President and Professor of Mathematics, The Russell College, Lebanon, Va.

A man has a park in the form of a parabolic segment cut off by a chord making an angle $\frac{1}{4}\pi$ with the axis. Within the park is a right angled triangular flower plat with one vertex at the center of gravity of the segment, the other vertex at the lower extremity of the chord, and the right angle on the diameter bisecting the chord. The park contains 30